X100/701

NATIONAL QUALIFICATIONS 2005 FRIDAY, 20 MAY 1.00 PM - 4.00 PM

MATHEMATICS ADVANCED HIGHER

Read carefully

- 1. Calculators may be used in this paper.
- 2. Candidates should answer all questions.
- 3. Full credit will be given only where the solution contains appropriate working.





4

3

4

4

Answer all the questions.

- 1. (a) Given $f(x) = x^3 \tan 2x$, where $0 < x < \frac{\pi}{4}$, obtain f'(x).
 - (b) For $y = \frac{1+x^2}{1+x}$, where $x \neq -1$, determine $\frac{dy}{dx}$ in simplified form.
- 2. Given the equation $2y^2 2xy 4y + x^2 = 0$ of a curve, obtain the x-coordinate of each point at which the curve has a horizontal tangent.
- Write down the Maclaurin expansion of e^x as far as the term in x⁴.
 Deduce the Maclaurin expansion of e^{x²} as far as the term in x⁴.

Hence, or otherwise, find the Maclaurin expansion of e^{x+x^2} as far as the term in x^4 .

4. The sum, S(n), of the first n terms of a sequence, u_1, u_2, u_3, \ldots is given by $S(n) = 8n - n^2, n \ge 1$.

Calculate the values of u_1 , u_2 , u_3 and state what type of sequence it is.

Obtain a formula for u_n in terms of n, simplifying your answer.

- 5. Use the substitution u = 1 + x to evaluate $\int_0^3 \frac{x}{\sqrt{1+x}} dx$.
- **6.** Use Gaussian elimination to solve the system of equations below when $\lambda \neq 2$:

$$x + y + 2z = 1$$

$$2x + \lambda y + z = 0$$

$$3x + 3y + 9z = 5.$$

Explain what happens when $\lambda = 2$.

7. Given the matrix $A = \begin{pmatrix} 0 & 4 & 2 \\ 1 & 0 & 1 \\ -1 & -2 & -3 \end{pmatrix}$, show that $A^2 + A = kI$ for some

constant k, where I is the 3×3 unit matrix.

Obtain the values of p and q for which $A^{-1} = pA + qI$.

8. The equations of two planes are x - 4y + 2z = 1 and x - y - z = -5. By letting z = t, or otherwise, obtain parametric equations for the line of intersection of the planes.

Show that this line lies in the plane with equation

$$x + 2y - 4z = -11.$$


9. Given the equation $z + 2i\overline{z} = 8 + 7i$, express z in the form a + ib.

10. Prove by induction that, for all positive integers n,

$$\sum_{r=1}^{n} \frac{1}{r(r+1)(r+2)} = \frac{1}{4} - \frac{1}{2(n+1)(n+2)}.$$

State the value of
$$\lim_{n\to\infty} \sum_{r=1}^{n} \frac{1}{r(r+1)(r+2)}$$
.

The diagram shows part of the graph of $y = \frac{x^3}{x-2}$, $x \ne 2$. 11.



Write down the equation of the vertical asymptote.

1

4

- Find the coordinates of the stationary points of the graph of $y = \frac{x^3}{x^2}$. (b)
- Write down the coordinates of the stationary points of the graph of $y = \left| \frac{x^3}{x - 2} \right| + 1.$ 2
- 12. Let $z = \cos \theta + i \sin \theta$.
 - Use the binomial expansion to express z^4 in the form u + iv, where u and v are expressions involving $\sin \theta$ and $\cos \theta$.
- 3
- Use de Moivre's theorem to write down a second expression for z^4 . (b)
- 1

Using the results of (a) and (b), show that (c)

$$\frac{\cos 4\theta}{\cos^2 \theta} = p \cos^2 \theta + q \sec^2 \theta + r, \text{ where } -\frac{\pi}{2} < \theta < \frac{\pi}{2},$$

stating the values of p, q and r.

6

[Turn over for Questions 13, 14 and 15 on Page four

Marks

4

13. Express
$$\frac{1}{x^3 + x}$$
 in partial fractions.

Obtain a formula for I(k), where $I(k) = \int_1^k \frac{1}{x^3 + x} dx$, expressing it in the form $\frac{a}{b}$, where a and b depend on k.

Write down an expression for $e^{I(k)}$ and obtain the value of $\lim_{k\to\infty} e^{I(k)}$.

14. Obtain the general solution of the differential equation

$$\frac{d^2y}{dx^2} - 3\frac{dy}{dx} + 2y = 20\sin x.$$
 7

Hence find the particular solution for which y = 0 and $\frac{dy}{dx} = 0$ when x = 0.

15. (a) Given
$$f(x) = \sqrt{\sin x}$$
, where $0 < x < \pi$, obtain $f'(x)$.

- (b) If, in general, $f(x) = \sqrt{g(x)}$, where g(x) > 0, show that $f'(x) = \frac{g'(x)}{k\sqrt{g(x)}}$, stating the value of k.

 Hence, or otherwise, find $\int \frac{x}{\sqrt{1-x^2}} dx$.
- (c) Use integration by parts and the result of (b) to evaluate

$$\int_0^{1/2} \sin^{-1} x \ dx.$$

[END OF QUESTION PAPER]